

Assessment of Shale-Oil Resources of the Sirte Basin Province, Libya, 2019

Using a geology-based assessment methodology, the U.S. Geological Survey estimated undiscovered, technically recoverable mean resources of 23.7 billion barrels of shale oil and 23 trillion cubic feet of associated gas in the onshore part of the Sirte Basin Province of Libya.

Introduction

The U.S. Geological Survey (USGS) quantitatively assessed the potential for undiscovered, technically recoverable continuous oil and gas resources in the onshore part of the Sirte Basin Province of Libya (fig. 1). The Sirte Basin Province encompasses a series of northwest–southeast trending horsts and grabens (or troughs) that resulted from multiple phases of rifting and subsidence beginning in the Triassic and continuing into the Neogene (Montgomery, 1994; Abadi and others, 2008; Badalini and others, 2009). Rifting and subsidence in the Sirte Basin Province were the result of the diachronous opening of the Atlantic and Tethys Oceans, differential movement of the African and Apulian plates, and transtensional and transpressional motion along the regional strike-slip faults in northernmost Africa (Fiduk, 2009; Hassan and Kendall, 2014). Major petroleum source rocks include the Lower Cretaceous Nubian shale and the Upper Cretaceous Sirte, Rachmat, and Etel Shales (Hassan and Kendall, 2014; Abualkhir, 2016). Subsidence in the Paleogene generally resulted in burial sufficient to thermally mature all source rocks into the oil-generation window (Gumati and Schamel, 1988; Futyan and Jawzi, 1996). All petroleum source rocks were interpreted to be of sufficient thickness and quality for each to be a potential target for horizontal drilling and were assessed separately. Lack of data prevented the delineation of shale-gas assessment units, although gas is present in a limited area of the basin (Hassan and Kendall, 2014).

Total Petroleum Systems and Assessment Units

The USGS defined four total petroleum systems (TPSs) and four assessment units (AUs) within these systems: (1) Sirte Shale TPS with the Sirte Shale Oil AU, (2) Rachmat Shale TPS with the Rachmat Shale Oil AU, (3) Etel Shale TPS with the Etel Shale Oil AU, and (4) Nubian Shale TPS with the Nubian Shale Oil AU. The Sirte shale is present and best developed as a petroleum source rock in all major troughs (Hassan and Kendall, 2014); the Rachmat and Etel Shales are in the Maradah, Ajdabiya, and Hameimat troughs, and the Nubian shale is in the Zallah and Hameimat troughs (fig. 1). The assessment unit areas could change as more information becomes available on the geochemistry of each petroleum source rock. Assessment input data are summarized in table 1.

The geologic model for these four TPSs is for oil to have been generated from each of the organic-rich shales in the Paleogene with peak oil generation possibly in the late Paleogene to Neogene (Hassan and Kendall, 2014). Some portion of the oil was partially retained within each of the shales following migration into conventional traps. The Sirte Shale contains Type II marine organic matter, has total organic carbon (TOC) contents of as much as 8 weight percent, hydrogen indices as high as 600 milligrams of hydrocarbon per gram of TOC, and shale thickness of as much as 600 meters (m; Hassan and Kendall, 2014). The Rachmat Shale contains Type II marine organic matter, has TOC contents of as much as 4 weight percent, and shale thickness of as much as 700 m. The Etel Shale contains Type II marine organic matter, has TOC contents of as much as 6.5 weight percent, and has shale thickness of as much as 300 m. The Lower Cretaceous Nubian shale is a lacustrine shale with Type I organic matter, has TOC contents of as much as 9 weight percent, and is as much as 500 m thick (Hassan and Kendall, 2014). Shales of

the Cenomanian–Turonian Eagle Ford Group were used as a partial production analog for the Sirte, Rachmat, and Etel Shales (Whidden and others, 2018), and lacustrine shale of the Uteland Butte member of the Green River Formation (nomenclature from Osmond, 1992) was used as a partial analog for the Nubian shale (Johnson and others, 2015).

Undiscovered Resources Summary

The USGS quantitatively assessed undiscovered continuous oil and gas resources in four assessment units (table 2) in the Sirte Basin Province of Libya. For undiscovered, technically recoverable shale-oil and associated gas resources, the mean totals are 23,706 million barrels of oil (MMBO), or 23.7 billion barrels of oil, with an F95–F5 fractile range from 5,505 to 50,652 MMBO; 23,033 billion cubic feet of associated gas (BCFG), or 23 trillion cubic feet of gas, with an F95–F5 fractile range from 5,000 to 51,464 BCFG; and 332 million barrels of natural gas liquids (MMBNGL) with an F95–F5 fractile range from 71 to 756 MMBNGL.




Figure 1. Map showing four continuous assessment units (AUs) in the Sirte Basin Province of Libya.

Table 1. Key input data for four continuous assessment units in the Sirte Basin Province of Libya.

[AU, assessment unit; %, percent; EUR, estimated ultimate recovery per well; MMBO, million barrels of oil. Well drainage area, success ratio, and EUR are defined partly using U.S. shale-oil analogs. The average EUR input is the minimum, median, maximum, and calculated mean. Shading indicates not applicable]

Assessment input data— Continuous AUs	Sirte Shale Oil AU				Rachmat Shale Oil AU			
	Minimum	Mode	Maximum	Calculated mean	Minimum	Mode	Maximum	Calculated mean
Potential production area of AU (acres)	1,000	9,500,000	24,987,000	11,496,000	1,000	7,272,500	14,545,000	7,272,833
Average drainage area of wells (acres)	60	100	140	100	60	100	140	100
Area untested in AU (%)	100	100	100	100	100	100	100	100
Success ratio (%)	10	50	90	50	10	50	90	50
Average EUR (MMBO)	0.06	0.15	0.28	0.155	0.06	0.15	0.28	0.155
AU probability	1.0				1.0			
Assessment input data— Continuous AUs	Etel Shale Oil AU				Nubian Shale Oil AU			
	Minimum	Mode	Maximum	Calculated mean	Minimum	Mode	Maximum	Calculated mean
Potential production area of AU (acres)	1,000	7,272,500	14,545,000	7,272,833	1,000	7,419,500	14,839,000	7,419,833
Average drainage area of wells (acres)	60	100	140	100	60	100	140	100
Area untested in AU (%)	100	100	100	100	100	100	100	100
Success ratio (%)	10	50	90	50	10	50	90	50
Average EUR (MMBO)	0.06	0.15	0.28	0.155	0.06	0.085	0.15	0.088
AU probability	1.0				1.0			

Table 2. Results for four continuous assessment units in the Sirte Basin Province of Libya.

[MMBO, million barrels of oil; BCFG, billion cubic feet of gas; NGL, natural gas liquids; MMBNGL, million barrels of natural gas liquids. Results shown are fully risked estimates. F95 represents a 95-percent chance of at least the amount tabulated; other fractiles are defined similarly. Fractiles are additive under the assumption of perfect positive correlation. Shading indicates not applicable]

Total petroleum systems and assessment units (AUs)	AU probability	Accu- mulation type	Total undiscovered resources											
			Oil (MMBO)				Gas (BCFG)				NGL (MMBNGL)			
			F95	F50	F5	Mean	F95	F50	F5	Mean	F95	F50	F5	Mean
Sirte Shale Total Petroleum System														
Sirte Shale Oil AU	1.0	Oil	2,026	7,914	19,681	9,009	1,893	7,724	20,457	8,994	28	114	312	135
Rachmat Shale Total Petroleum System														
Rachmat Shale Oil AU	1.0	Oil	1,327	5,090	12,057	5,680	1,235	4,937	12,617	5,675	18	73	192	85
Etel Shale Total Petroleum System														
Etel Shale Oil AU	1.0	Oil	1,357	5,132	12,034	5,707	1,276	4,999	12,609	5,714	19	74	192	86
Nubian Shale Total Petroleum System														
Nubian Shale Oil AU	1.0	Oil	795	3,010	6,880	3,310	596	2,336	5,781	2,650	6	23	60	26
Total undiscovered continuous resources			5,505	21,146	50,652	23,706	5,000	19,996	51,464	23,033	71	284	756	332

References Cited

Abadi, A.M., van Wees, J.-D., van Dijk, P.M., and Cloetingh, S.A.P.L., 2008, Tectonics and subsidence evolution of the Sirt Basin, Libya: American Association of Petroleum Geologists Bulletin, v. 92, no. 8, p. 993–1027.

Abualkhir, E.A., 2016, Libya and the great challenges of overcoming difficulties to exploring and producing shale gas, and tight reservoirs (shale oil) potential: American Association of Petroleum Geologists, Search and Discovery Article No. 10850, 22 p., accessed March 26, 2019, at http://www.searchanddiscovery.com/documents/2016/10850abualkhir/ndx_abualkhir.pdf.

Badalini, G., Redfern, J., and Carr, I.D., 2009, A synthesis of current understanding of the structural evolution of North Africa: Journal of Petroleum Geology, v. 25, no. 3, p. 249–258.

Fiduk, J.C., 2009, Evaporites, petroleum exploration, and the Cenozoic evolution of the Libyan shelf margin, central North Africa: Marine and Petroleum Geology, v. 26, no. 8, p. 1513–1527.

Futyan, A., and Jawzi, A.H., 1996, The hydrocarbon habitat of the oil and gas fields of North Africa with emphasis on the Sirt Basin, in Salem, M.J., El-Hawati, A.S., and Sbata, A.M., eds., Geology of the Sirt Basin, v. 2: Amsterdam, Elsevier, p. 287–308.

Gumati, Y.D., and Schamel, S., 1988, Thermal maturation history of the Sirte Basin, Libya: Journal of Petroleum Geology, v. 11, no. 2, p. 205–218.

Hassan, H.S., and Kendall, C.C.G., 2014, Hydrocarbon provinces of Libya—A petroleum system study, chap. 4 of Marlow, L., and Yose, L., eds., Petroleum systems of the Tethyan region: American Association of Petroleum Geologists Memoir No. 106, p. 101–141.

Johnson, R.C., Birdwell, J.E., Mercier, T.J., Brownfield, M.E., Charpentier, R.R., Klett, T.R., Leathers, H.M., Schenk, C.J., and Tennyson, M.E., 2015, Assessment of undiscovered oil and gas resources in the Uteland Butte Member of the Eocene Green River Formation, Uinta Basin, Utah: U.S. Geological Survey Fact Sheet 2015–3052, 2 p.

Montgomery, S., 1994, Sirte Basin, north-central Libya—Prospects for the future: Petroleum Frontiers, v. 11, no. 1, 92 p.

Osmond, J.C., 1992, Greater Natural Buttes gas field, Uintah County, Utah, in Fouch, T.D., Nuccio, V.F., and Chidsey, T.C., Jr., eds., Hydrocarbon and mineral resources of the Uinta Basin, Utah and Colorado: Utah Geological Association Guidebook No. 20, p. 143–163.

Whidden, K.J., Pitman, J.K., Pearson, O.N., Paxton, S.T., Kinney, S.A., Gianoutsos, N.J., Schenk, C.J., Leathers-Miller, H.M., Birdwell, J.E., Brownfield, M.E., Burke, L.A., Dubiel, R.F., French, K.L., Gaswirth, S.B., Haines, S.S., Le, P.A., Marra, K.R., Mercier, T.J., Tennyson, M.E., and Woodall, C.A., 2018, Assessment of undiscovered oil and gas resources in the Eagle Ford Group and associated Cenomanian–Turonian strata, U.S. Gulf Coast, Texas, 2018: U.S. Geological Survey Fact Sheet 2018–3033, 4 p.

Differences in the spelling of the Sirte (Sirt) Basin reflect the individual author's usage.

Sirte Basin Province Assessment Team

Christopher J. Schenk, Tracey J. Mercier, Cheryl A. Woodall, Phuong A. Le, Janet K. Pitman, Ronald M. Drake II, Michael E. Brownfield, Stephanie B. Gaswirth, and Thomas M. Finn

For More Information

Assessment results are also available at the USGS Energy Resources Program website at <https://energy.usgs.gov>.